

Local-energy density functionals for an N-dimensional electronic system in a magnetic field

B.P. van Zyl

Department of Physics, Queen's University,
Kingston, Ontario, Canada K7L 3N6

(February 1, 2008)

We present a general approach for the construction of the exact local-energy density functionals for a uniform N-dimensional electronic system in a magnetic field. For *arbitrary* dimension, we obtain explicit expressions for the matter, kinetic, and exchange density functionals. In the zero-field limit, we recover the usual N-dimensional Thomas-Fermi theory. As an application of our results, we develop a current-density-functional theory, in the spirit of the Thomas-Fermi-Dirac approximation, for an inhomogeneous many-electron system in a magnetic field.

I. INTRODUCTION

It is now well established that a density-functional approach to the ground state and dynamical properties of many-electron systems is a viable alternative to the usual wavefunction approach involving the solution of the single-particle Schrödinger equation. The proof by Vignale and Rasolt¹ (VR) that the scalar and vector potentials characterizing a many-electron system are uniquely determined by the single-particle electron and current densities (the so-called current-density-functional theory CDFT) has placed density-functional theory on a rigorous footing for the description of many-electron systems involving electric and magnetic fields.

In this work, we present a general approach for the construction of the local-energy density functionals for an N-dimensional electronic system in a homogeneous magnetic field. Unlike the Kohn-Sham single-particle scheme, in which only the exchange-correlation functional is unknown, this statistical formalism requires the knowledge of both the kinetic and exchange energy functionals in the presence of electric and magnetic fields. Our approach is to consider the exact canonical density matrix for a noninteracting N-dimensional electron gas in a homogeneous magnetic field. The single-particle density matrix is obtained via the inverse Laplace transform of the canonical density matrix, as originally formulated by March and Murray². The diagonal component of the single-particle density matrix yields the exact density distribution of the system. The exact kinetic and exchange energy-density functionals may also be explicitly evaluated from the single-particle density matrix. Although these functionals are obtained for a homogeneous system, an application of the local-density approximation (LDA), in the spirit of Thomas-Fermi-Dirac theory³ (TFD), allows these functionals to be also used in inhomogeneous

systems.

Our paper is organized as follows. In Sec. II, we define the N-dimensional canonical density matrix for a noninteracting electron gas in a homogeneous magnetic field, and discuss some of the mathematical properties of the associated N-dimensional single-particle density matrix. Then, in Sec. III, we consider the evaluation of the kinetic and exchange energy-density functionals for arbitrary dimension and magnetic field strength. Explicit forms for the kinetic and exchange energy-densities are obtained for three-and-two-dimensional systems. In Sec. IV, we present an outline for the construction of a current-density-functional theory based on the TFD approximation in two-and-three-dimensions, making use of the density functionals obtained in Sec. III. Finally, in Sec. V we offer our concluding remarks.

II. DENSITY MATRIX OF A N-DIMENSIONAL HOMOGENEOUS ELECTRON GAS IN A UNIFORM MAGNETIC FIELD

Long ago, Sondheimer and Wilson⁴ considered the evaluation of the diamagnetism of free electrons. The central tool in their calculation was the determination of the canonical density matrix, which in 3-dimensions (3D) is given by (including spin degeneracy)

$$C(\mathbf{r}, \mathbf{r}'; \beta) = \left(\frac{m}{2\pi\hbar^2\beta} \right)^{3/2} \frac{\hbar\omega\beta}{\sinh(\hbar\omega\beta/2)} \times \exp \left\{ -\frac{m}{2\hbar^2\beta} [i\hbar\omega\beta(x'y - y'x) + \frac{\hbar\beta}{2} \coth(\hbar\omega\beta/2)((x - x')^2 + (y - y')^2) + (z - z')^2] \right\}, \quad (1)$$

where the symmetric gauge⁵ has been used and $\omega = eB/mc$. For purely formal reasons, we extend the definition of $C(\mathbf{r}, \mathbf{r}'; \beta)$ to N-dimensions in the following way

$$C(\mathbf{r}, \mathbf{r}'; \beta) = \left(\frac{m}{2\pi\hbar^2\beta} \right)^{N/2} \frac{\hbar\omega\beta}{\sinh(\hbar\omega\beta/2)} \times \exp \left\{ -\frac{m}{2\hbar^2\beta} [i\hbar\omega\beta(x'y - y'x) + \frac{\hbar\omega\beta}{2} \coth(\hbar\omega\beta/2)((x - x')^2(y - y')^2)] \right\}$$

$$\times \prod_{i=1}^{N-2} \exp \left(-\frac{m}{2\hbar^2\beta} (z_i - z'_i)^2 \right), \quad (2)$$

with $(N = 2, 3, \dots)$. The single-particle density matrix in N -dimensions is then obtained via the usual inverse Laplace transform relation⁶

$$\rho(\mathbf{r}, \mathbf{r}') = \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{d\beta}{2\pi i} \frac{e^{\beta\epsilon_f}}{\beta} C(\mathbf{r}, \mathbf{r}'; \beta), \quad (3)$$

with ϵ_f being the Fermi energy, and the contour chosen to be closed in the left-half of the β -plane (see Fig. 1). We defer an exact treatment of Eq. (3) until the next section.




FIG. 1. (a) The contour to be used when d is integral, and (b) when d is half-integral.

The electron number density ρ (spin-averaged, zero temperature) is immediately given by taking the trace of the density matrix in Eq. (3), viz.,

$$\rho^{(d)}(\gamma) = 2 \left(\frac{m\omega}{4\pi\hbar} \right)^d \mathcal{F}^{(d)}(\gamma), \quad (4)$$

with

$$\mathcal{F}^{(d)}(\gamma) \equiv \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{ds}{2\pi i} \frac{e^{2\gamma s}}{s^d \sinh(s)}, \quad (5)$$

and for notational convenience, we have defined $s \equiv \hbar\omega\beta/2$, $\gamma \equiv \epsilon_f/(\hbar\omega)$, and $d = N/2$. Depending on the dimension of the space, d can be either integral or half-integral.

In the case that d is half-integral (i.e., odd- N), the inverse Laplace transform defined by $\mathcal{F}^{(d)}$ *always* contains a branch point in the complex s -plane at $s = 0$. Therefore, care must be taken when evaluating $\mathcal{F}^{(d)}(\gamma)$ at the branch point. The contour chosen for the evaluation of Eq. (5) is shown in Fig. 1b, where a cut has been introduced along the negative real axis, and the large arc does not pass through any of the poles of the integrand. Following the method of Sondheimer and Wilson,⁴ we find that the integral in Eq. (5) can be written as a Riemann zeta function plus the sum of the residues at the poles $s_k = \pm k\pi i$

$$\mathcal{F}^{(d)}(\gamma) \equiv \int_{\sigma} \frac{ds}{2\pi i} \frac{e^{2\gamma s}}{s^d \sinh(s)} + \sum_k \text{Res} \left(\frac{e^{2\gamma s_k}}{s_k^d \sinh(s_k)} \right), \quad (6)$$

where σ denotes the contour FEDC in Fig. 1b. The first integral in Eq. (6) can be separated into two parts by using an asymptotic expansion of the integrand at the branch point

$$\begin{aligned} \int_{\sigma} \frac{ds}{2\pi i} \frac{e^{2\gamma s}}{s^d \sinh(s)} &= \int_{\sigma} \frac{ds}{2\pi i} e^{2\gamma s} \sum_{k=0}^{d+\frac{1}{2}} \frac{B_k}{s^{d+1-k}} \\ &- \int_{\sigma} \frac{ds}{2\pi i} e^{2\gamma s} \left(\sum_{k=0}^{d+\frac{1}{2}} \frac{B_k}{s^{d+1-k}} - \frac{1}{s^d \sinh(s)} \right). \end{aligned} \quad (7)$$

where

$$B_k = \frac{1}{k!} \frac{\partial^k}{\partial s^k} \left(\frac{s}{\sinh(s)} \right)_{s=0}. \quad (8)$$

The first integral is evaluated by means of Hankel's formula for the gamma function, viz.,⁶

$$\int_{\sigma} \frac{ds}{2\pi i} e^{2\gamma s} s^{-z} = \frac{(2\gamma)^{z-1}}{\Gamma(z)}, \quad (9)$$

and the second integral can be cast as real since the part contributed by the circle around the origin (see Fig. 1b) is zero. The sum of the residues is readily computed, and we finally have

$$\begin{aligned} \mathcal{F}^{(d)}(\gamma) &= \sum_{k=0}^{d+\frac{1}{2}} \frac{B_k (2\gamma)^{d-k}}{\Gamma(d+1-k)} + \frac{1}{\pi} \mathcal{G}^{(d)}(\gamma) \\ &(-1)^{d+\frac{1}{2}} \sum_{k=1}^{\infty} \frac{2(-1)^k}{(\pi k)^d} \cos(2\pi\gamma k + \pi d/2) \end{aligned} \quad (10)$$

where

$$\mathcal{G}^{(d)}(\gamma) = \int_0^{\infty} dy e^{2\gamma y} \left(\sum_{k=0}^{d+\frac{1}{2}} \frac{B_k}{y^{d+1-k}} - \frac{1}{y^d \sinh(y)} \right). \quad (11)$$

The first term in Eq. (10) gives the TF expression for the number density in a homogeneous magnetic field. The second and third terms are purely quantum-mechanical, and are the origin of the de Haas-van Alphen effect.⁷ These oscillatory terms are clearly a result of the exact treatment of the canonical density matrix, and are not obtained within the usual TF approximation.⁸

Integral values of d allow for an exact treatment of Eq. (5). Using the contour shown in Fig. 1a, $\mathcal{F}^{(d)}(\gamma)$ can be expressed in the form

$$\mathcal{F}^{(d)}(\gamma) = \sum_k \text{Res} \left(\frac{e^{2\gamma s_k}}{s_k^d \sinh(s_k)} \right), \quad (12)$$

where $s_k = k\pi i$ ($k = 0, \pm 1, \pm 2, \dots$). The $k = 0$ contribution represents the usual TF approximation, whereas

the terms associated with the poles at $\pm k\pi i$ are responsible for the quantum-mechanical de Haas-van Alphen effect. As in the half-integral case, $\mathcal{F}^{(d)}(\gamma)$ for even- N will be the sum of steady terms ($s_k = 0$) plus oscillatory terms ($s_k = \pm k\pi i$):

$$\mathcal{F}^{(d)} = A_{k=0}^{(d)} + (-1)^d \sum_{k=1}^{\infty} \frac{2(-1)^k}{(\pi k)^d} \cos(2\pi\gamma k + \pi d/2) \quad (13)$$

where the $A_{k=0}^{(d)}$ terms are given by

$$A_{k=0}^{(d)} = \frac{1}{d!} \frac{\partial^d}{\partial s^d} \left(\frac{e^{2\gamma s}}{\sinh(s)} \right)_{s=0}. \quad (14)$$

In the limit $\omega \rightarrow 0$, the asymptotic behaviour of $\mathcal{F}^{(d)}(\gamma)$ for arbitrary d is given by⁹

$$\mathcal{F}^{(d)}(\gamma) = \frac{(2\gamma)^d}{\Gamma(d+1)}, \quad (15)$$

and Eq. (4) reduces to the N -dimensional TF theory, viz.,^{8,10}

$$\rho_{\text{TF}}^{(d)} = \left(\frac{m}{2\pi\hbar^2} \right)^d \frac{2}{\Gamma(d+1)} \varepsilon_f^d. \quad (16)$$

For large magnetic fields, the general behaviour of $\mathcal{F}^{(d)}(\gamma)$ for half-integral d is difficult to obtain using the present treatment. Once again, the cause of this difficulty lies in the branch point at $s = 0$ of the complex s -plane. We can circumvent this problem by writing Eq. (5) in the following form¹¹

$$\mathcal{F}^{(d)}(\gamma) = 2 \sum_{k=0}^{\infty} \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{ds}{2\pi i} \frac{e^{(2\gamma s - 2k-1)s}}{s^d}, \quad (17)$$

where we have used the series expansion

$$\frac{1}{\sinh(s)} = 2 \sum_{k=0}^{\infty} e^{-(2k+1)s}, \quad (18)$$

for the $\sinh(s)$ function. Only those terms satisfying $2k+1 < 2\gamma$ survive the integration in Eq. (17), and since $\max[2\gamma] = 2$, we have

$$\mathcal{F}^{(d)}(\gamma) = \frac{2}{\Gamma(d)} (2\gamma - 1)^{d-1}. \quad (19)$$

Equations (4) and (19) provide the high-magnetic field electronic number density for arbitrary values of d .

III. ENERGY-DENSITY FUNCTIONALS

The kinetic and exchange energy functionals can both be calculated from the single-particle density matrix,

given by Eq. (3). However, in the present context, a simplification for the evaluation of the kinetic energy density is possible. Our first step is to notice that the Heavyside function may be represented as the inverse Laplace transform of the exponential function:¹²

$$\theta(\varepsilon_f - \hat{H}) = \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{d\beta}{2\pi i} \frac{e^{\beta(\varepsilon_f - \hat{H})}}{\beta}. \quad (20)$$

The kinetic energy density ε_{kin} may then be expressed as

$$\begin{aligned} \varepsilon_{\text{kin}} &= \langle \mathbf{r} | \hat{H} \theta(\varepsilon_f - \hat{H}) | \mathbf{r} \rangle \\ &= \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{d\beta}{2\pi i} \frac{\langle \mathbf{r} | \hat{H} e^{\beta(\varepsilon_f - \hat{H})} | \mathbf{r} \rangle}{\beta} \\ &= - \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{d\beta}{2\pi i} \frac{e^{\beta\varepsilon_f}}{\beta} \frac{\partial C(\mathbf{r}, \mathbf{r}; \beta)}{\partial \beta}. \end{aligned} \quad (21)$$

For arbitrary dimension, ε_{kin} takes the form

$$\begin{aligned} \varepsilon_{\text{kin}}^{(d)} &= \left(\frac{m}{4\pi} \right)^d \hbar^{d-1} \omega^{d+1} \int_{\epsilon-i\infty}^{\epsilon+i\infty} \frac{ds}{2\pi i} e^{2\gamma s} \times \\ &\quad \left[\frac{(d-1)}{s^{d+1} \sinh(s)} + \frac{\cosh(s)}{s^d \sinh(s)^2} \right] \\ &= \left(\frac{m}{4\pi} \right)^d \hbar^{1-d} \omega^{d+1} \left[2\gamma \mathcal{F}^{(d)}(\gamma) - \mathcal{F}^{(d+1)}(\gamma) \right] \\ &= \left(\frac{4\pi\hbar^2}{m} \right) \left(\frac{\rho^{(d)}}{2} \right)^{\frac{d+1}{d}} H^{(d)}(\gamma), \end{aligned} \quad (22)$$

where

$$H^{(d)}(\gamma) = \frac{2\gamma}{[\mathcal{F}^{(d)}]^{1/d}} - \frac{[\mathcal{F}^{(d+1)}]}{[\mathcal{F}^{(d)}]^{(d+1)/d}}. \quad (23)$$

Equation (22) is valid for arbitrary magnetic field strength.

In the limit of zero magnetic field, the asymptotic behaviour of $H^{(d)}(\gamma)$ for arbitrary d can be deduced from Eq. (15), viz.,

$$H^{(d)}(\gamma) = \frac{\Gamma(d+2)[\Gamma(d+1)]^{\frac{1}{d}} - [\Gamma(d+1)]^{\frac{d+1}{d}}}{\Gamma(d+2)}. \quad (24)$$

Thus, using the above value of $H^{(d)}(\gamma)$ in Eq. (22), we recover the kinetic energy density of the N -dimensional TF theory.^{8,10} As the dimension of the space goes to infinity, we obtain

$$H^{(d)}(\gamma) = 1, \quad (25)$$

whereby the kinetic energy density becomes

$$\varepsilon_{\text{kin}}^{(d)} = 4\pi \left(\frac{\hbar^2}{2m} \right) \rho^{(d)}. \quad (26)$$

Substituting Eq. (19) into Eq. (23) gives the high-field limit of $H^{(d)}(\gamma)$, viz.,

$$H^{(d)}(\gamma) = \frac{4\gamma\Gamma(d)\frac{1}{d}\Gamma(d+1)(2\gamma-1)^{d-1}}{\Gamma(d+1)2^{\frac{d+1}{d}}(2\gamma-1)^{\frac{d^2-1}{d}}} - \frac{2(2\gamma-1)^d\Gamma(d)^{\frac{d+1}{d}}}{\Gamma(d+1)2^{\frac{d+1}{d}}(2\gamma-1)^{\frac{d^2-1}{d}}} \quad (27)$$

In the limit $d \rightarrow \infty$, we obtain

$$H^{(d)}(\gamma) = 2\gamma, \quad (28)$$

and the kinetic energy behaves as

$$\varepsilon_{\text{kin}}^{(d)} = 8\pi\gamma \left(\frac{\hbar^2}{2m} \right) \rho^{(d)} \simeq 8\pi \left(\frac{\hbar^2}{2m} \right) \rho^{(d)}. \quad (29)$$

The other density functional of interest is the N-dimensional exchange energy ε_{ex} . According to the usual prescription, ε_{ex} is given by³

$$\varepsilon_{\text{ex}} = -\frac{e^2}{4} \int d\mathbf{r}' \frac{|\rho(\mathbf{r}, \mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|}, \quad (30)$$

where the integration is understood to be over the N-dimensional volume. The exchange energy density has a much richer s -plane topology than the kinetic energy density because of the essential singularities arising from the $\coth(s)$ term in Eq. (2). These essential singularities make it difficult (if not impossible) to obtain a useful, closed analytical form for the N-dimensional exchange energy for *arbitrary* magnetic fields. Nevertheless, some analytical progress can be made if we make use of the following expansion:⁶

$$\frac{\exp(-\alpha \coth(s))}{\sinh(s)} = 2 \sum_{n=0}^{\infty} L_n(2\alpha) e^{-\alpha} e^{-(2n+1)s}, \quad (31)$$

where $L_n(\alpha) \equiv L_n^{(0)}(\alpha)$ is the Laguerre polynomial. It is then easy to show that the N-dimensional single-particle density matrix has the form

$$\rho(\mathbf{r}, \mathbf{r}') = 4 \left(\frac{m\omega}{4\pi\hbar} \right)^d e^{-\frac{m\omega}{2\hbar}(x'y-y'x)} \sum_{n=0}^{\infty} L_n(2\alpha) e^{-\alpha} \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \frac{ds}{2\pi i} \frac{e^{(2\gamma-2n-1)s} e^{-\delta/s}}{s^d}, \quad (32)$$

where we have defined

$$\alpha = \frac{m\omega}{4\hbar} |\mathbf{r} - \mathbf{r}'|^2 \quad ; \quad \delta = \frac{m\omega}{4\hbar} \sum_{i=1}^{2(d-1)} (z_i - z'_i)^2. \quad (33)$$

Equation (32) in Eq. (30) yields, in principle, the N-dimensional exchange energy for arbitrary magnetic field. Of course, the comments made in Sec. II with regard to the electron number density also apply to Eq. (32).

Below, we illustrate the utility of our formalism through specific examples in the physically relevant 3D and 2D systems.

The electron number density in three-dimensions can be obtained from Eqs. (4) and (10):

$$\rho^{(3/2)}(\gamma) = \frac{1}{4} \left(\frac{m\omega}{\pi\hbar} \right)^{3/2} \left[\frac{4}{3\sqrt{\pi}} (2\gamma)^{3/2} - \frac{1}{6\sqrt{2\pi\gamma}} + \frac{1}{\pi} \mathcal{G}^{(3/2)}(\gamma) + \sum_{k=1}^{\infty} \frac{2}{\pi^{3/2}} \frac{(-1)^k}{k^{3/2}} \cos \left(2k\pi\gamma + \frac{3\pi}{4} \right) \right], \quad (34)$$

where $\mathcal{G}^{(3/2)}(\gamma)$ is given by Eq. (11). In the zero field limit, we can make use of Eq. (16) to get

$$\rho^{(3/2)}(\gamma) = \frac{1}{3\pi^2} \left(\frac{2m}{\hbar^2} \right)^{3/2} \varepsilon_f^{3/2}, \quad (35)$$

and in the high-field limit, Eqs. (4) and (19) yield,

$$\rho^{(3/2)}(\gamma) = \left(\frac{2m^3\omega^2}{\pi^4\hbar^4} \right)^{1/2} (\varepsilon_f - \hbar\omega/2)^{1/2}. \quad (36)$$

Both of these limiting cases are in full agreement with the results expected for a 3D homogeneous electron gas.³

The kinetic energy density is most easily obtained by using the second form of Eq. (22) with $d = 3/2$, viz.,

$$\varepsilon_{\text{kin}} = \left(\frac{m}{4\pi} \right)^{3/2} \hbar^{-1/2} \omega^{5/2} \left[2\gamma \mathcal{F}^{(3/2)} - \mathcal{F}^{(5/2)} \right]. \quad (37)$$

The functions $\mathcal{F}^{(3/2)}(\gamma)$ and $\mathcal{F}^{(5/2)}(\gamma)$ are readily obtained from Eqs. (10), and (11), and therefore

$$\begin{aligned} \varepsilon_{\text{kin}} = & \left(\frac{m}{4\pi} \right)^{3/2} \hbar^{-1/2} \omega^{5/2} \left[\frac{4(2\gamma)^{5/2}}{5\sqrt{\pi}} + \frac{(2\gamma)^{1/2}}{6\sqrt{\pi}} \right. \\ & + \frac{2\gamma}{\pi} \mathcal{G}^{(3/2)}(\gamma) - \frac{1}{\pi} \mathcal{G}^{(5/2)}(\gamma) \\ & + \sum_{k=1}^{\infty} \frac{4\gamma(-1)^{(k+1)}}{(\pi k)^{3/2}} \sin(2k\pi\gamma + \frac{\pi}{4}) \\ & \left. + \sum_{k=1}^{\infty} \frac{2(-1)^k}{(\pi k)^{5/2}} \sin(2k\pi\gamma - \frac{\pi}{4}) \right]. \quad (38) \end{aligned}$$

The last four terms in Eq. (38) are again quantum effects and are the origin of the de Haas-van Alphen effect.

Using Eq. (24) in Eq. (22) yields the kinetic energy density in the zero-field limit

$$\varepsilon_{\text{kin}} = \frac{3}{5} (3\pi^2)^{2/3} \left(\frac{\hbar^2}{2m} \right) \rho^{5/3}, \quad (39)$$

and in the high-field regime, Eq. (27) in Eq. (22) gives

$$\varepsilon_{\text{kin}} = \frac{\hbar\omega}{2} \rho + \frac{2\pi^4\hbar^4}{3m^3\omega^2} \rho^3. \quad (40)$$

Equation (39) is the well known TF result for a degenerate noninteracting 3D electron gas,³ while Eq. (40)

agrees with the result of Yonei *et al.*,¹² and Banerjee *et al.*¹³ Note that the functional form of the kinetic energy density is very different in the low and high-magnetic field limits. In particular, the ρ^3 term appearing in Eq. (40) is a reflection of the one dimensional nature of the system along the field direction.

In order to evaluate the exchange energy density, we require explicit expressions for the single-particle density matrix. In principle, Eq. (32) provides such an expression. However, it is rather difficult to obtain the exchange energy for arbitrary magnetic field in 3D,¹⁴ so we focus here only on the the zero and high-field limits.

In the zero-field limit, we set $\sinh(s) \approx s$, and $\cosh(s) \approx s^{-1}$. Substituting these expressions into Eq. (2), and performing the inverse Laplace transform, we obtain

$$\rho(\mathbf{r}, \mathbf{r}') = \frac{\sqrt{\varepsilon_f}}{a\sqrt{\pi}} \cosh(2\sqrt{a\varepsilon_f}) - \frac{1}{2\sqrt{a^3\pi}} \sinh(2\sqrt{a\varepsilon_f}) , \quad (41)$$

where $a \equiv (m/2\pi\hbar^2) |\mathbf{r} - \mathbf{r}'|^2$. This expression leads back to Eq. (35) upon taking the limit $\mathbf{r} \rightarrow \mathbf{r}'$. Equation (30) can now be used to yield the exchange energy functional. This is a straightforward, but tedious calculation, and since it only leads back to the well know expression for a 3D free-electron gas, we simply quote the result here³

$$\varepsilon_{\text{ex}} = -\frac{3}{4} \left(\frac{3}{\pi} \right)^{1/3} e^2 \rho^{4/3} . \quad (42)$$

The high-field limit for the single-particle density matrix is obtained by replacing $\sinh(s) \approx 1/2\exp(-s)$, and $\cosh(s) \approx 1$ in Eq. (2). After performing the inverse Laplace transform of the canonical density matrix, we obtain

$$\begin{aligned} \rho(\mathbf{r}, \mathbf{r}') &= \frac{2\hbar\omega}{\sqrt{2\pi^2}} \left(\frac{m}{2\pi\hbar^2} \right)^{3/2} \frac{1}{|z - z'|} \\ &\times \sin(\sqrt{2(\varepsilon_f - \hbar\omega/2)} |z - z'|) \\ &\times \exp\left(-i\frac{m\omega}{2\hbar}(x'y - y'x)\right) \\ &\times \exp\left(-\frac{m\omega}{4\hbar}((x - x')^2 + (y - y')^2)\right) . \end{aligned} \quad (43)$$

(The high-field limits of the number density can be obtained from Eq. (43) by taking the limit $\mathbf{r} \rightarrow \mathbf{r}'$). The exchange energy density can now be calculated by substituting Eq. (43) into Eq. (30) and transforming over to cylindrical coordinates:

$$\begin{aligned} \varepsilon_{\text{ex}} &= -\frac{4e^2}{(2\pi)^3} \left(\frac{m\omega}{\hbar} \right)^2 \int_0^\infty dz z^{-2} \sin^2 \left[\left(\frac{4\pi^4 \hbar^4 \rho^2}{m^3 \omega^2} \right)^{1/2} z \right] \\ &\times \int_0^\infty dr r (r^2 + z^2)^{-1/2} \exp\left(-\frac{m\omega}{2\hbar} r^2\right) . \end{aligned} \quad (44)$$

Fortunately, Danz and Glasser¹⁴ have already considered a similar calculation, and after several manipulations in accordance with their work, we obtain

$$\begin{aligned} \varepsilon_{\text{ex}} &= -\frac{e^2}{4\pi^3} \left(\frac{m\omega}{\hbar} \right)^2 \left[p \int_0^\infty dx x \ln(x^2) \exp(-px^2) \right. \\ &\quad - p \int_0^\infty dx x \ln(x^2 + 1) \exp(-px^2) \\ &\quad \left. + 2p \int_0^\infty dx \exp(-px^2) \tan^{-1}(x^{-1}) \right] , \end{aligned} \quad (45)$$

where

$$p = \frac{8\pi^4 \hbar^3}{m^3 \omega^3} \rho^2 . \quad (46)$$

The asymptotic behaviour of Eq. (45) in the low-density limit is given by

$$\varepsilon_{\text{ex}} \approx \left(\frac{e^2}{8\pi^3} \right) \left(\frac{m\omega}{\hbar} \right)^2 p [\ln(p) - (3 - C)] , \quad (47)$$

where C is Euler's constant. In the high-density regime (where the semiclassical TF theory is applicable), Eq. (45) has the form

$$\varepsilon_{\text{ex}} \approx -\left(\frac{e^2}{8\pi^3} \right) \left(\frac{m\omega}{\hbar} \right)^2 p^{1/2} . \quad (48)$$

Similar to the kinetic energy density functional, the behaviour of the exchange energy density is markedly different in the low and high magnetic field limits.

B. N=2

Since d is integral, we can make use of Eq. (13) to obtain the electron number density in terms of elementary functions, viz.,

$$\begin{aligned} \rho^{(1)}(\gamma) &= \left(\frac{m\omega}{2\pi\hbar} \right) \mathcal{F}^{(1)}(\gamma) \\ &= \left(\frac{m\omega}{2\pi\hbar} \right) \left[2\gamma + \sum_{k=1}^\infty \frac{2(-1)^k}{\pi k} \sin(2\pi k\gamma) \right] , \end{aligned} \quad (49)$$

The low and high-field limits of the number density are given by

$$\rho^{(1)}(\gamma) = \frac{m\varepsilon_f}{\pi\hbar^2} , \quad (50)$$

and

$$\rho^{(1)}(\gamma) = \frac{m\omega}{\pi\hbar} , \quad (51)$$

respectively.

The kinetic energy density can also be expressed in terms of elementary functions, viz.,

$$\begin{aligned}
\varepsilon_{\text{kin}} &= \left(\frac{m\omega^2}{4\pi} \right) \left[2\gamma \mathcal{F}^{(1)}(\gamma) - \mathcal{F}^{(2)}(\gamma) \right] \\
&= \left(\frac{m\omega^2}{4\pi} \right) \left[2\gamma^2 + \frac{1}{6} + \sum_{k=1}^{\infty} \frac{4\gamma(-1)^k}{\pi k} \sin(2\pi k\gamma) \right. \\
&\quad \left. + \sum_{k=1}^{\infty} \frac{2(-1)^k}{(\pi k)^2} \cos(2\pi k\gamma) \right] \\
&= \left(\frac{\hbar^2}{m} \right) \pi \left[\rho^{(1)}(\gamma) \right]^2 H^{(1)}(\gamma) \\
&= \left(\frac{\hbar^2}{2m} \right) \pi \left[\rho^{(1)}(\gamma) \right]^2, \tag{52}
\end{aligned}$$

where we have used the fact that $H^{(1)}(\gamma) = 1/2 \forall \gamma$. Therefore, in remarkable contrast to the 3D case, the 2D kinetic energy density has *exactly* the same functional form regardless of the magnetic field strength; all of the de Haas-van Alphen type behaviour is buried in the $\rho^{(1)}(\gamma)$ dependence. This result is peculiar to 2D, since in all other dimensions, $H^{(d)}(\gamma)$ is generally a function of ω . For example, in four-dimensions, we have from Eq. (24)

$$\varepsilon_{\text{kin}} = \pi \left(\frac{\hbar^2}{2m} \right) \frac{8}{3} \rho^{3/2}, \tag{53}$$

in the zero-field limit, and from Eq. (27)

$$\varepsilon_{\text{kin}} = \frac{\pi^2 \hbar^3}{m^2 \omega} \rho^2 + \frac{\hbar \omega}{2} \rho, \tag{54}$$

in the high-field limit.

The exchange energy is calculated along the same lines as Sec. III A. The single-particle density matrix for arbitrary magnetic field is obtained by setting $d = 1$ in Eq. (32), and is given by

$$\begin{aligned}
\rho(\mathbf{r}, \mathbf{r}') &= \frac{m\omega}{\pi \hbar} \exp \left(\frac{-im\omega}{2\hbar} (x'y - y'x) \right) \times \\
&\quad \exp \left(-\frac{m\omega}{4\hbar} |\mathbf{r} - \mathbf{r}'|^2 \right) \times \\
&\quad L_{n_f}^{(1)} \left(\frac{m\omega}{2\hbar} |\mathbf{r} - \mathbf{r}'|^2 \right), \tag{55}
\end{aligned}$$

where we have summed only up to the Fermi level n_f , and used the finite summation relation⁶

$$\sum_{n=0}^m L_n^{(k)}(x) = L_m^{(k+1)}(x). \tag{56}$$

The exchange energy in 2D can be evaluated from Eq. (30), provided that we make use of the identity⁶

$$\left[L_n^{(1)}(x) \right]^2 = \frac{\Gamma(2+n)}{n!} \sum_{k=0}^{\infty} \frac{(2n-2k)!(2k)! L_{2k}^{(2)}(2x)}{\Gamma(2-k) (n-k)!}. \tag{57}$$

Upon substituting Eq. (55) into Eq. (30), we obtain after some tedious algebra

$$\begin{aligned}
\varepsilon_{\text{ex}} &= -\frac{e^2}{\pi} \left(\frac{m\omega}{2\hbar} \right)^{3/2} n_f \sum_{k=0}^{n_f} \\
&\quad \frac{\Gamma(n_f + 2 + k) \Gamma(2k + 1/2)}{\Gamma(n_f + 1 - k) \Gamma(2k + 2) \Gamma(k + 2) \Gamma(k + 1)} \times \\
&\quad F[-(n_f - k), (2k + 1/2); (2k + 2); 2], \tag{58}
\end{aligned}$$

where $F(a, b; c; d)$ denotes the hypergeometric series.⁶ The limiting forms of Eq. (58) in the zero and high-field limits can be made more transparent by first examining at the limiting forms of $\rho(\mathbf{r}, \mathbf{r}')$ in Eq. (3).

In the zero-field limit, we have

$$\rho(\mathbf{r}, \mathbf{r}') = \left(\frac{1}{\pi} \right) \frac{1}{|\mathbf{r} - \mathbf{r}'|} k_f J_1(k_f |\mathbf{r} - \mathbf{r}'|), \tag{59}$$

where the Fermi wavevector $k_f = (2m\varepsilon_f/\hbar^2)^{1/2}$. Once again, Eq. (50) is recovered by taking the limit $\mathbf{r} \rightarrow \mathbf{r}'$ in Eq. (59). The exchange energy in this limit is given by an application of Eq. (30), viz.,

$$\begin{aligned}
\varepsilon_{\text{ex}} &= -\frac{2}{3} e^2 \frac{1}{\pi^2} k_f^3 \\
&= -\frac{4}{3} \sqrt{\frac{2}{\pi}} e^2 \rho^{3/2}. \tag{60}
\end{aligned}$$

This is the usual 2D free-electron gas result. In the high field limit, the single-particle density matrix takes the form

$$\begin{aligned}
\rho(\mathbf{r}, \mathbf{r}') &= \frac{m\omega}{\pi \hbar} \exp \left(-i \frac{m\omega}{2\pi \hbar} (x'y - y'x) \right) \\
&\quad \times \exp \left(-\frac{m\omega}{4\pi \hbar} ((x - x')^2 + (y - y')^2) \right), \tag{61}
\end{aligned}$$

and upon using Eq. (30), we find the simple result

$$\varepsilon_{\text{ex}} = -\frac{\pi e^2}{2\sqrt{2}} \rho^{3/2}. \tag{62}$$

Equations (60), and (61) should be contrasted with Eqs. (42), and (45). Obviously, the key difference between the two sets of functionals is the universal form assumed by the exchange energy in 2D. Specifically, the exchange energy functional in 2D is explicitly independent of γ in limit of strong magnetic fields.

IV. CURRENT-DENSITY FUNCTIONAL THEORY IN STRONG MAGNETIC FIELDS: THE TFD APPROXIMATION

In this section, we take a pedestrian approach to the construction of a current density functional theory applicable to inhomogeneous electronic systems in strong magnetic fields. The foundations of the CDFT formalism have already been provided by VR¹, and we invite the interested reader to refer to this work for rigorous proofs and details.

Consider an *inhomogeneous* electronic system in a magnetic field. Within the TFD approximation, we assume that the non-uniform electronic system can be locally approximated as a uniform electron gas characterized by the density $\rho(\mathbf{r})$, local Fermi wavevector $k_f(\mathbf{r})$, and canonical (or “conjugate”) current density $\mathbf{j}_p(\mathbf{r})$ (see below). Thus, a local approximation to the energy functional is obtained by summing over all N-dimensional volume elements, treating each N-dimensional volume element as independent; this is the local-density approximation (LDA) in which the density is assumed to vary slowly on the scale of the magnetic length $\ell_B = (\hbar c/eB)^{1/2}$.

The local approximation to the total ground state energy functional for an N-dimensional many-electron system characterized by a scalar potential $v(\mathbf{r})$ and vector potential $\mathbf{A}(\mathbf{r})$ is given by

$$E_{\text{TFD}} = \int d\mathbf{r} \varepsilon[\rho(\mathbf{r}), |\boldsymbol{\nu}(\mathbf{r})|] + \frac{m}{2} \int d\mathbf{r} \frac{|\mathbf{j}(\mathbf{r})|^2}{\rho(\mathbf{r})} + E_H[\rho(\mathbf{r})] + \int d\mathbf{r} \rho(\mathbf{r})v(\mathbf{r}), \quad (63)$$

where E_H is the nonlocal Hartree contribution, and $\varepsilon = \varepsilon_{\text{kin}} + \varepsilon_{\text{ex}}$ is the total (gauge-invariant) energy density of the uniform electron gas in a fictitious magnetic field, defined by the vorticity $\boldsymbol{\nu}(\mathbf{r})$

$$\boldsymbol{\nu}(\mathbf{r}) = \nabla \times \frac{\mathbf{j}_p(\mathbf{r})}{\rho(\mathbf{r})}. \quad (64)$$

The physical orbital current density is given by

$$\mathbf{j}(\mathbf{r}) = \mathbf{j}_p(\mathbf{r}) + \frac{e}{mc} \rho(\mathbf{r}) \mathbf{A}(\mathbf{r}), \quad (65)$$

where

$$\mathbf{j}_p(\mathbf{r}) \equiv \frac{\hbar}{m} \text{Im}[\nabla \rho(\mathbf{r}, \mathbf{r}')|_{r=r'}]. \quad (66)$$

The rationale behind considering the total energy ε in a fictitious magnetic field is as follows: For a given local value of $\mathbf{j}_p$, we construct a fictitious vector potential $\mathbf{A}_\nu(\mathbf{r})$ which produces $\mathbf{j}_p$ in a uniform electron gas of density ρ . The potential $\mathbf{A}_\nu(\mathbf{r})$ couples only to the current and density. Since we must have $\mathbf{j} = 0$ everywhere within a uniform electron gas,¹⁵ we find

$$\mathbf{A}_\nu(\mathbf{r}) = -\frac{mc}{e} \frac{\mathbf{j}_p}{\rho(\mathbf{r})}. \quad (67)$$

The magnetic field associated with $\mathbf{A}_\nu(\mathbf{r})$ is nothing more than the vorticity defined in Eq. (64). The relationship between the fictitious magnetic field and the true magnetic field is given by

$$\boldsymbol{\nu}(\mathbf{r}) = \boldsymbol{\omega} - \nabla \times \frac{\mathbf{j}(\mathbf{r})}{\rho(\mathbf{r})}. \quad (68)$$

Therefore, the functional ε is simply constructed by considering the Euler equation for a homogeneous electron gas in a fictitious magnetic field $\boldsymbol{\nu}(\mathbf{r})$, viz.,

$$\frac{\partial \varepsilon[\rho(\mathbf{r}), |\boldsymbol{\nu}|]}{\partial \rho(\mathbf{r})} = \mu = \mu_{\text{kin}} + \mu_{\text{ex}} \quad (69)$$

where the chemical potential μ now consists of kinetic plus exchange contributions to the free electron gas in the fictitious magnetic field.

The minimization of Eq. (63) is done first with respect to $\mathbf{j}(\mathbf{r})$ (or equivalently $\mathbf{j}_p$, since ρ and $\mathbf{A}$ are held constant). This leads to an Euler-Lagrange equation for the equilibrium orbital current density, which may be cast in terms of the density $\rho(\mathbf{r})$. A substitution of the resulting expression for the orbital current density into Eq. (63) yields a functional, E_{TFD} , solely in terms of $\rho(\mathbf{r})$. A minimization of this functional gives the Euler-Lagrange equation for the ground state density distribution of the many-electron system.

Below, we treat separately the 3D and 2D systems within the CDFT, restricting our attention to the case of strong magnetic fields. The energy-density functionals obtained in Secs. III A and III B now find relevance in the construction of the total energy density $\varepsilon = \varepsilon_{\text{kin}} + \varepsilon_{\text{ex}}$.

A. N=3

The total energy density ε is obtained from Eq. (69), with

$$\mu = \frac{\hbar |\boldsymbol{\nu}|}{2} + \frac{2\hbar^4 \pi^4}{m^3 |\boldsymbol{\nu}|^2} \rho^2 - e^2 \left(\frac{m}{8\pi^2 \hbar} \right)^{\frac{1}{2}} |\boldsymbol{\nu}|^{\frac{1}{2}}, \quad (70)$$

where the last term is the exchange contribution to the chemical potential. Using Eq. (70), we immediately obtain

$$\begin{aligned} \varepsilon[\rho, |\boldsymbol{\nu}|] &= \frac{\hbar}{2} |\boldsymbol{\nu}| \rho + \frac{2\pi^4 \hbar^4}{3m^3 |\boldsymbol{\nu}|^2} \rho^3 \\ &\quad - e^2 \left(\frac{m}{8\pi^2 \hbar} \right)^{1/2} |\boldsymbol{\nu}|^{1/2} \rho. \end{aligned} \quad (71)$$

Notice that Eq. (71) is formally equivalent to replacing $\boldsymbol{\omega} \rightarrow |\boldsymbol{\nu}|$ in Eqs. (40) and (45) (e.g., the total energy density in the *fictitious* magnetic field). As discussed above, a variation of the E_{TFD} with respect to $\mathbf{j}$ gives

$$\begin{aligned} \mathbf{j} &= \frac{1}{m} \nabla \times \frac{\partial \varepsilon}{\partial \boldsymbol{\nu}} \\ &= \left(\frac{\hbar}{2m} - \frac{4\pi^4 \hbar^4}{m^4} \frac{\rho^2}{|\boldsymbol{\nu}|^3} - \frac{e^2}{2} \left(\frac{m}{8\pi^2 \hbar} \right)^{1/2} \right) \nabla \rho \times \hat{\mathbf{z}}. \end{aligned} \quad (72)$$

In the limit of very strong fields, we obtain

$$\mathbf{j} = \frac{\hbar}{2m} \nabla \rho \times \hat{\mathbf{z}}, \quad (73)$$

which is exactly the result of Vignale and Skudlarski.¹⁶ The vorticity can now be developed by using Eq. (72)

in Eq. (68). Inserting the resulting expression for the vorticity into Eq. (71) gives all of the necessary components for the complete characterization of the ground state energy functional, Eq. (63).

One can easily improve on the above TFD theory by including gradient corrections (the so-called von Weizsäcker corrections¹⁷) in the kinetic energy density. The explicit inclusion of such gradient terms would take into account the magnetic-induced anisotropy of the density along, and perpendicular to the field direction, thereby giving a more realistic treatment of the system in a strong magnetic field. Such a calculation has already been performed by Mazzolo *et al.*,¹⁸ but without the exchange interaction. Their final result for the total energy functional is valid within the Thomas-Fermi-von Weizsäcker approximation, and is given by

$$E_{\text{TFvW}} = \int d\mathbf{r} \left(\frac{\hbar\omega}{2} \rho + \frac{2\pi^4 \hbar^4}{3m^3 \omega^2} \rho^3 - \frac{\hbar^2}{8m} \frac{|\nabla_{\perp} \rho|^2}{\rho} + \frac{3\pi^4 \hbar^5}{m^4 \omega^3} \rho |\nabla_{\perp} \rho|^2 - 8m \left(\frac{\pi^4 \hbar^4}{m^4 \omega^3} \right)^2 \rho^3 |\nabla_{\perp} \rho|^2 - \frac{\hbar^2}{24m} \frac{|\nabla_{\parallel} \rho|^2}{\rho} \right) + \int d\mathbf{r} \rho v(\mathbf{r}), \quad (74)$$

where $\nabla_{\perp}$ and $\nabla_{\parallel}$ are the gradients perpendicular and parallel to the magnetic field direction respectively. Our formalism offers an improvement over Eq. (74) by including the effects of the exchange interaction.

B. N=2

The determination of the total energy density ε in 2D is formally the same as in the 3D case. However, the simple substitution $\omega \rightarrow |\boldsymbol{\nu}|$ in Eqs. (52) and (61) for the present scenario is quite ambiguous. Obviously, this is because the energy-density functionals for $N = 2$ are explicitly independent of the magnetic field in the strong-field regime. So, unlike the 3D case, we have to be a little more careful when constructing the 2D total energy density ε .

First, we define the chemical potential for the homogeneous 2D electron gas in the fictitious magnetic field:

$$\mu = \frac{\hbar |\boldsymbol{\nu}|}{2} - \frac{\pi e^2}{2\sqrt{2}} \left(\frac{m}{2\pi\hbar} \right)^{1/2} |\boldsymbol{\nu}|^{1/2} \quad (75)$$

An application of Eq. (69) yields

$$\varepsilon[\rho, |\boldsymbol{\nu}|] = \frac{\hbar |\boldsymbol{\nu}|}{2} \rho - \frac{\pi e^2}{2\sqrt{2}} \left(\frac{m}{2\pi\hbar} \right)^{1/2} |\boldsymbol{\nu}|^{1/2} \rho. \quad (76)$$

If we now use Eq. (51) with $\omega \rightarrow |\boldsymbol{\nu}|$, Eq. (76) may be cast in terms of the density ρ , viz.,

$$\varepsilon[\rho, |\boldsymbol{\nu}|] = \pi \left(\frac{\hbar^2}{2m} \right) \rho^2 - \frac{\pi e^2}{2\sqrt{2}} \rho^{3/2}. \quad (77)$$

Recall that in 3D, the magnetic-TFD theory is readily distinguished from the non-magnetic theory because the local 3D energy functionals enjoy very different behaviour in the low and high-field limits. Conversely, the local 2D energy-density functionals have no such signature.

The current density is obtained with the aid of Eq. (76), viz.,

$$\begin{aligned} \mathbf{j} &= \frac{1}{m} \nabla \times \frac{\partial \varepsilon}{\partial \boldsymbol{\nu}} \\ &= \left(\frac{\hbar}{2m} - \frac{e^2}{8m} \left(\frac{m}{2\pi\hbar |\boldsymbol{\nu}|} \right)^{\frac{1}{2}} \right) \nabla \rho \times \hat{\mathbf{z}} \\ &\simeq \left(\frac{\hbar}{2m} - \frac{e^2}{8m} \left(\frac{m}{2\pi\hbar \omega} \right)^{\frac{1}{2}} \right) \nabla \rho \times \hat{\mathbf{z}}. \end{aligned} \quad (78)$$

In the limit of very strong fields we again recover Eq. (73). MacDonald and Girvin¹⁹ have shown some time ago that Eq. (73) is a special case of a powerful identity expressing the single-particle density matrix in terms of the density. Indeed, for a 2D electron gas in a uniform magnetic field, Eq. (73) is the *exact* orbital current density distribution when only the lowest Landau level is occupied.

The recent interest in the optical, and electronic properties of the 2D electron gas in a strong magnetic field (e.g., quantum-hall effect²⁰, 2D Wigner crystallization²¹, edge reconstruction in quantum dots²²) provides adequate motivation for the explicit calculation of Eq. (63) in two-dimensions.

The vorticity is determined from Eq. (78) and Eq. (64), viz.,

$$\begin{aligned} |\boldsymbol{\nu}(\mathbf{r})| &= \omega - \left(\frac{\hbar}{2m} - \frac{\pi e^2}{4\sqrt{2}m} \left(\frac{m}{2\pi\hbar\omega} \right)^{1/2} \right) \\ &\quad \times \frac{|\nabla_{\perp} \rho|^2}{\rho^2}, \end{aligned} \quad (79)$$

where terms of the form $\nabla_{\perp}^2 \rho$ are ignored, as they eventually integrate to zero. Using this expansion in Eq. (76), and with Eq. (78) for the current, we obtain

$$\begin{aligned} \varepsilon + \frac{m}{2} \frac{|\mathbf{j}|^2}{\rho} &= \frac{\hbar\omega}{2} \rho - \frac{\pi e^2}{2\sqrt{2}} \left(\frac{m}{\pi\hbar} \right)^{\frac{1}{2}} \omega^{1/2} \rho \\ &\quad - \left[\frac{\hbar^2}{8m} - \frac{e^2}{8} \left(\frac{\pi\hbar}{2m\omega} \right)^{\frac{1}{2}} + \frac{\pi e^4}{64\hbar\omega} \right] \\ &\quad \times \frac{|\nabla_{\perp} \rho|^2}{\rho}. \end{aligned} \quad (80)$$

Using Eq. (80) in Eq. (63), we finally obtain for the total energy functional, E_{TFD}

$$E_{\text{TFD}} = \int d\mathbf{r} \left[\frac{\hbar\omega}{2} \rho - \frac{\pi e^2}{2\sqrt{2}} \left(\frac{m}{\pi\hbar} \right)^{\frac{1}{2}} \omega^{1/2} \rho \right]$$

$$- \left(\frac{\hbar^2}{8m} - \frac{e^2}{8} \left(\frac{\pi\hbar}{2m\omega} \right)^{\frac{1}{2}} + \frac{\pi e^4}{64\hbar\omega} \right) \frac{|\nabla_{\perp}\rho|^2}{\rho} + v(\mathbf{r})\rho(\mathbf{r}) + \frac{e^2}{2} \int d\mathbf{r}' \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \quad (81)$$

Notice that the first two terms have the same form as the usual zero-field TFD energy functional (the kinetic term is *exactly* the same as the zero-field TF term). Extending Eq. (81) to include gradient corrections in the kinetic energy is not possible because the analogous 2D von Weizsäcker term vanishes identically.⁸ It is interesting to note, however, that the current density naturally provides a von Weizsäcker-like (vW) gradient correction, viz.,

$$E_{\text{vW}} = -\frac{\hbar^2}{2m} \int d\mathbf{r} \frac{|\nabla_{\perp}\rho|^2}{4\rho}, \quad (82)$$

with a coefficient that is 1/9-th the magnitude of the 3D, zero-field TFDvW theory.⁸ It is well known that the von Weizsäcker term provides the smooth exponential fall-off for the density distribution as $\mathbf{r} \rightarrow \infty$.

Equation (81) is applicable to any smooth external confining potential $v(\mathbf{r})$. For example, choosing $v(\mathbf{r}) = 1/2m\omega_0 r^2$ in Eq. (81) would provide a scheme for calculating the ground state density distribution of a parabolically confined quantum dot in a strong magnetic field.

V. CONCLUSIONS

We have presented a general approach for the calculation of the exact local-density functionals for an N-dimensional electronic system in a magnetic field. We have found that 2D systems enjoy a preferred status over other dimensions, in that the form of the 2D kinetic and exchange energy functionals are explicitly independent of the magnetic field. We have also provided an outline for the construction of a CDFT for 3D and 2D systems, which illustrates how the local-density functionals obtained in this work may be used to describe non-uniform electronic systems in a strong magnetic field. The results presented here may be of use in a wide variety of problems of current physical interest. In particular, our results for 2D systems should be of relevance to researchers interested in density-functional methods for low-dimensional electronic systems.

Although the present work has focused on spin-averaged densities at zero temperature, an extension to include spin and finite temperatures is straightforward.

VI. ACKNOWLEDGEMENTS

It is a pleasure to thank Dr. R. K. Bhaduri for stimulating discussions. The hospitality of the Department of

Physics and Astronomy at McMaster University, Hamilton, Canada, is also gratefully acknowledged. Financial support for this work was provided by the NSERC of Canada and Queen's University, Kingston, Canada.

-
- ¹ G. Vignale, and M. Rasolt, Phys. Rev. Lett. **59**, 2360, 1987; Phys. Rev. B **37** 10685 (1988).
 - ² N. H. March, and A. M. Murray, Phys. Rev. **120**, 830 (1960).
 - ³ R. M. Dreizler, and E.K.U Gross, *Density Functional Theory: An Approach To The Quantum Many-Body Problem*, (Springer-Verlag, Berlin, 1990).
 - ⁴ E. H. Sondheimer, and A. H. Wilson, Proc. R. Soc. London Ser. **A210**, 173 (1951).
 - ⁵ The choice of gauge is arbitrary, because the total energy-functional must be gauge invariant.
 - ⁶ I. S. Gradshteyn and I.M. Ryzhik, *Table of Integrals* (Academic, New York, 1980).
 - ⁷ C. Kittel, *Introduction to Solid State Physics* (Wiley, New York, 1996), p. 257.
 - ⁸ M. Brack and R. K. Bhaduri, *Semiclassical Physics*, Frontiers in Physics Vol. 96 (Addison-Wesley, Reading, 1997), Ch. 4.
 - ⁹ Of course, care must be still be exercised at the branch point, $s = 0$, when evaluating the large γ behaviour of $\mathcal{F}^{(d)}(\gamma)$.
 - ¹⁰ G. F. Kventzel and J. Katriel, Phys. Rev. A **24**, 2299 (1981).
 - ¹¹ D. Shoenberg, Proc. R. Soc. London Ser. **A170**, 341 (1939).
 - ¹² K. Yonei and T. Motomochi, Jour. Phys. Soc. Jap. **59**, 3571 (1990).
 - ¹³ B. Banerjee, D. H. Constantinescu and P. Rehák, Phys. Rev. D **10**, 2384 (1974).
 - ¹⁴ R. W. Danz and M. L. Glasser, Phys. Rev. B **4**, 94 (1971).
 - ¹⁵ L. D. Landau, Z. Phys. **64**, 629 (1930).
 - ¹⁶ P. Skudlarski and G. Vignale, Phys. Rev. B **48**, 8547 (1993).
 - ¹⁷ C. F. v. Weizsäcker, Z. Phys. **96**, 431 (1935).
 - ¹⁸ A. Mazzolo, E. L. Pollock and G. Zérah, Phys. Lett. A **209**, 123 (1995).
 - ¹⁹ A. H. MacDonald and S. M. Girvin, Phys. Rev. B **38**, 6295 (1988).
 - ²⁰ *The Quantum Hall Effect*, edited by R. E. Prange and S. M. Girvin (Springer-Verlag, New York, 1987).
 - ²¹ E. P. Wigner, Phys. Rev. **46**, 1002 (1934).
 - ²² Los Alamos preprint server: cond-mat/9904067